

Applicant : Jens Chr. Jensen
Serial No. : 09/874,198
Filed : June 4, 2001
Page : 3

A [REDACTED] Docket No.: 09011-002002

REMARKS

Applicants hereby submit that the enclosures fulfill the requirements under 37 C.F.R. §1.821-1.825. The amendments in the specification merely insert the paper copy of the Sequence Listing and sequence identifiers in the specification. No new matter has been added.


Attached hereto is a marked-up version of the changes made to the specification by the current amendment.

Please apply any charges or credits to Deposit Account No. 06-1050.

Respectfully submitted,

Date: _____

1/3/02



John W. Freeman, Esq.
Reg. No. 29,066

Fish & Richardson P.C.
225 Franklin Street
Boston, Massachusetts 02110-2804
Telephone: (617) 542-5070
Facsimile: (617) 542-8906

20310520.doc

09011-002002-06142850

"Version With Markings to Show Changes Made"

In the specification:

Paragraph beginning at page 15, line 4, has been amended as follows:

Figure 2 shows the sequence alignment²¹ of the amino acid sequences of MASP-2 (clone phl-4; amino acid residues 16-686 of SEQ ID NO:2), MASP-1^{17,22} (SEQ ID NO:6), C1r^{23,24} (SEQ ID NO:7) and C1s^{25,26} (SEQ ID NO:8).

Paragraph beginning at page 15, line 14, has been amended as follows:

Figure 6 shows the cDNA sequence and deduced amino acid sequence of MASP-2 (SEQ ID NOs:3 and 2, respectively).

Paragraph beginning at page 46, line 10, has been amended as follows:

The liver is the primary site of synthesis of C1r, C1s, and MASP-1. Thus, RNA from liver was used as template for RT-PCR with primers deduced from the obtained peptide sequences. First strand synthesis of cDNA was carried out with 1.3 μ g human liver RNA using a First-Strand cDNA Synthesis Kit (Pharmacia). PCR was performed on this cDNA using degenerate sense and antisense primers derived from the amino acid sequences EYANDQER (SEQ ID NO:4) and KPFTGFEA (SEQ ID NO:5), respectively. The PCR program consisted of 1 cycle with annealing at 50EC; 1 cycle with annealing at 55EC, and 33 cycles with annealing at 60EC. The resulting 300 bp PCR product was cloned into the *E. coli* plasmid pCRII using the TA-cloning kit (InVitrogen) and the nucleotide sequence of the insert was determined.

09874198-09011-002002